arXiv:1302.3095v3 [math.GM] 9 Apr 2013

IMPROVING THREE-POINT ITERATIVE METHODS FOR SOLVING NONLINEAR EQUATIONS

FAYYAZ AHMAD AND D. GARCA-SENZ

Abstract. In this article, we report on sixth-order and seventh-order iterative methods for solving nonlinear equations. In particular sixth-order derivative-based and derivative-free iterative families are constructed in such a way that they comprise a wide class of sixth-order methods which were developed in the past years. Weighting functions are introduced to enhance the algorithmic efficiency whereas an appropriate parametric combination gives weight-age flexibility in between those weighting functions. The usage of weighting factors and weighting functions define a wide class of iterative schemes for solving nonlinear equations. The freedom to construct different parametric combinations as well as different forms of weighting functions makes the iterative schemes more accurate and flexible, It means that one can easily modify the scheme by changing weight functions and parametric combination.

1. Introduction

Nonlinear algebraic equations are at the heart of many problems of nonlinear science. However, it is not always possible to find exact analytical solutions of these nonlinear equations which very often have to be solved numerically. Iterative methods are numerical algorithms which, starting from an initial guess in the neighborhood of a root of the nonlinear equation, manage to refine it and achieve convergence to the true root after several iterations . We can divide them in two classes, namely memory-based and without-memory iterative methods. From the point of view of efficiency and stability, multipoint iterative methods are superior than single-point methods. Kung and Traub conjectured [13] that multipoint iterative methods without-memory have at most 2^n order of convergence for $n+1$ function evaluations per iteration. The further classification falls into derivative-free and non-derivative-free iterative methods. The multipoint iterative methods for nonlinear equations, which use derivatives, are efficient. The most famous derivative-based iterative method is Newton's method (**NM**) which is written as

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}, \qquad n = 0, 1, 2, 3, \cdots . \tag{1}$$

It is quadratically convergent in some neighborhood of the root α of $f(x) = 0$. The computational effort invested in the derivative of a function is not always comparable to the evaluation of the function itself [2, 3, 4, 5, 6, 7, 8]. Steffensen [1] gave a quadratically convergent derivative-free iterative method for nonlinear equations. The Steffensen's method (**SM**) is given by

$$\begin{cases} w_n &= x_n - \kappa f(x_n), \\ df(x_n) &= \frac{f(x_n) - f(w_n)}{\kappa f(x_n)} \\ x_{n+1} &= x_n - \frac{f(x_n)}{df(x_n)}. \end{cases} \tag{2}$$

Clearly, $df(x_n)$ is a finite difference approximation of $f'(x_n)$. The **SM** approximation merely replaces the derivative evaluation by a function evaluation. Actually both iterative method **NM** and **SM** achieve the optimal order of convergence 2, which is in agreement with the conjecture of Kung and Traub. The computational efficiency of an iterative method (IM) is frequently calculated by the Ostrowski-Traub's [14, 15] efficiency index

$$E(IM) = p^{1/d}, \tag{3}$$

where p is order of convergence and d is function evaluations per iteration. The optimal computational efficiency is defined by expression

$$E_{opt}^{(n)} = 2^{n/(n+1)}. \tag{4}$$

We now briefly review a number of three step sixth-order convergent methods. Many three steps sixth-order convergent methods use three function and one derivative evaluations. Sharma and Guha [9] proposed the following sixth-order convergent family by introducing one parameter (**SG**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= y_n - \frac{f(x_n)}{f(x_n) - 2f(y_n)} \frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} &= z_n - \frac{f(x_n) + af(y_n)}{f(x_n) + (a-2)f(y_n)} \frac{f(z_n)}{f'(x_n)}, \end{cases} \tag{5}$$

Key words and phrases. Non-linear equations, Iterative methods, Derivative free, Steffensen's method, Convergence order.

where $a \in \Re$. Neta [10] uni-parametric sixth-order family consists of three step (**NT1**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= y_n - \frac{f(x_n)+af(y_n)}{f(x_n)+(a-2)f(y_n)} \frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} &= z_n - \frac{f(x_n)-f(y_n)}{f(x_n)-3f(y_n)} \frac{f(z_n)}{f'(x_n)}, \end{cases} \tag{6}$$

where $a \in \Re$. In 2011, Neta [11] provided a more efficient sixth-order scheme (**NT2**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= y_n - \frac{f(y_n)}{f'(x_n)} \frac{1}{\left[1-\frac{f(y_n)}{f(x_n)}\right]^2}, \\ x_{n+1} &= z_n - \frac{f(z_n)}{f'(x_n)} \frac{1}{\left[1-\frac{f(y_n)}{f(x_n)}-\frac{f(z_n)}{f(x_n)}\right]^2}. \end{cases} \tag{7}$$

Chun and Ham [12] also presented a family of sixth-order iterative methods (**CH**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= y_n - \frac{f(x_n)}{f(x_n)-2f(y_n)} \frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} &= z_n - H(\mu_n) \frac{f(z_n)}{f'(x_n)}, \end{cases} \tag{8}$$

where real valued function $H(s)$ satisfies $H(0) = 1$, $H'(0) = 2$, and $\mu_n = f(y_n)/f(x_n)$. Grau et al. [16] developed the sixth-order (**GR**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= y_n - \frac{f(x_n)}{f(x_n)-2f(y_n)} \frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} &= z_n - \frac{f(x_n)}{f(x_n)-2f(y_n)} \frac{f(z_n)}{f'(x_n)}. \end{cases} \tag{9}$$

Alicia cordero et al. [17] constructed the following sixth-order scheme (**AL**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= x_n - \theta \frac{f(x_n)+f(y_n)}{f'(x_n)} - (1-\theta) \frac{f(x_n)}{f'(x_n)} \frac{f(x_n)}{f(x_n)-f(y_n)}, \\ x_{n+1} &= z_n - \frac{f(z_n)}{f[z_n,y_n]+f[z_n,x_n,x_n](z_n-y_n)}, \end{cases} \tag{10}$$

where $\theta \in \Re$.Finally, Sanjay K. Khattri et al. [21] presented a paper, Unification of sixth-order iterative methods, which covers all the features of sixth-order iterative methods (**SK1**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= y_n - \frac{f(y_n)}{f'(x_n)} \left(1 + \sum_{j=1}^{m} \left(\frac{f(y_n)}{f(x_n)}\right)^j a_j\right), \text{ where } a_1 = 2, \\ x_{n+1} &= z_n - \frac{f(z_n)}{f'(x_n)} \left(1 + \sum_{k=1}^{l} \left(\frac{\mu_1 f(y_n)+\mu_2 f(z_n)}{f(x_n)}\right)^k b_k\right), \text{ where } b_1 = 2/\mu_1, \end{cases} \tag{11}$$

where, $\{a_j,\, b_k,\, \mu_1,\, \mu_2\} \in \Re$, $m \in \mathbf{Z}^+$, and $l \in \mathbf{Z}^+$ are independent parameters. Sanjay K. Khattri et al. [18] also developed four parameter sixth-order derivative free family (**SK2**)

$$\begin{cases} y_n &= x_n - \kappa \frac{f(x_n)^2}{f(x_n)-f(x_n-\kappa f(x_n))}, \\ z_n &= y_n - \kappa \frac{f(x_n)f(y_n)}{f(x_n)-f(x_n-\kappa f(x_n))} \left[1 + \frac{f(y_n)}{f(x_n)} + \alpha \left(\frac{f(y_n)}{f(x_n)}\right)^2 + \frac{f(y_n)}{f(x_n-\kappa f(x_n))} + \beta \left(\frac{f(y_n)}{f(x_n-\kappa f(x_n))}\right)^2\right], \\ x_{n+1} &= z_n - \kappa \frac{f(x_n)f(z_n)}{f(x_n)-f(x_n-\kappa f(x_n))} \left[1 + \frac{f(y_n)}{f(x_n)} + \alpha \left(\frac{f(y_n)}{f(x_n)}\right)^2 + \frac{f(y_n)}{f(x_n-\kappa f(x_n))} + \beta \left(\frac{f(y_n)}{f(x_n-\kappa f(x_n))}\right)^2 + \eta \frac{f(z_n)}{f(y_n)}\right]. \end{cases} \tag{12}$$

In [19], R. Thukral constructed Steffensen-Six.one method (**TS1**) and Steffensen-six.two method (**TS2**), which are respectively

$$\begin{cases} w_n &= x_n + f(x_n), \\ y_n &= x_n - \left(\frac{f(x_n)^2}{f(w_n)-f(x_n)}\right), \\ z_n &= y_n - \left(1 - \lambda^{-1} \frac{f(y_n)}{f(w_n)}\right)^{-\lambda} \left(\frac{x_n-y_n}{f(x_n)-f(y_n)}\right) f(y_n), \\ x_{n+1} &= z_n - \left(1 - \lambda^{-1} \frac{f(y_n)}{f(w_n)}\right)^{-\lambda} \left(\frac{x_n-y_n}{f(x_n)-f(y_n)}\right) f(z_n), \end{cases} \tag{13}$$

and

$$\left\{\begin{aligned}
w_n &= x_n + f(x_n),\\
y_n &= x_n - \left(\frac{f(x_n)^2}{f(w_n)-f(x_n)}\right),\\
z_n &= y_n - \left(1-\lambda^{-1}\frac{f(y_n)}{f(w_n)}\right)^{-\lambda}\left(\frac{x_n-y_n}{f(x_n)-f(y_n)}\right)f(y_n),\\
x_{n+1} &= z_n - \left(\frac{z_n-y_n}{f(z_n)-f(y_n)}\right)f(z_n),
\end{aligned}\right. \tag{14}$$

where $\lambda \in \Re/\{0\}$. F. Soleymani et al. [20] discussed the following sixth-order derivative-free method (**FS1**)

$$\left\{\begin{aligned}
w_n &= x_n + f(x_n),\\
y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]},\\
z_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]}\left[1+\frac{f(y_n)}{f(x_n)}\left(1+2\frac{f(y_n)}{f(x_n)}\right)\right],\\
x_{n+1} &= z_n - \frac{f(z_n)}{f[y_n,z_n]}\left(1-\frac{1+f[x_n,w_n]}{f[x_n,w_n]}\frac{f(z_n)}{f(w_n)}\right),
\end{aligned}\right. \tag{15}$$

where $f[y_n,z_n] = \frac{f(z_n)-f(y_n)}{z_n-y_n}$, $f[x_n,w_n] = \frac{f(w_n)-f(x_n)}{w_n-x_n}$. F. Soleymani [22] also presented an other sixth-order derivative-free scheme (**FS2**)

$$\left\{\begin{aligned}
w_n &= x_n - \beta f(x_n),\\
y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]},\\
z_n &= y_n - \frac{f(y_n)}{f[w_n,y_n]},\\
x_{n+1} &= z_n - \frac{f(z_n)}{f[w_n,z_n]+f[z_n,y_n]-f[w_n,y_n]},
\end{aligned}\right. \tag{16}$$

where $\beta \in \Re/\{0\}$.In 2013, F. Soleymani [26] developed some efficient seventh-order derivative-free methods. First is (**FS3-1**)

$$\left\{\begin{aligned}
w_n &= x_n + f(x_n),\\
y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]},\\
z_n &= y_n - \frac{f(y_n)}{f[x_n,y_n]+f[y_n,w_n]-f[x_n,w_n]},\\
x_n &= z_n - \frac{f(z_n)}{f[x_n,z_n]}\left[1+\frac{f(y_n)}{f(w_n)}+\frac{f(z_n)}{f(y_n)}+\left\{\frac{2+f[x_n,w_n]}{(1+f[x_n,w_n])^2}\right\}\left(\frac{f(y_n)}{f(x_n)}\right)^2+\gamma\frac{f(z_n)}{f(x_n)}+\delta\frac{f(z_n)}{f(w_n)}\right],
\end{aligned}\right. \tag{17}$$

where γ and δ are real numbers. Second is (**FS3-2**)

$$\left\{\begin{aligned}
w_n &= x_n - f(x_n),\\
y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]},\\
z_n &= y_n - \frac{f(y_n)}{f[x_n,y_n]+f[y_n,w_n]-f[x_n,w_n]},\\
x_n &= z_n - \frac{f(z_n)}{f[x_n,z_n]}\left[1+\frac{f(y_n)}{f(w_n)}+\frac{f(z_n)}{f(y_n)}+\left\{\frac{2-f[x_n,w_n]}{(1-f[x_n,w_n])^2}\right\}\left(\frac{f(y_n)}{f(x_n)}\right)^2+\rho\frac{f(z_n)}{f(x_n)}+\tau\frac{f(z_n)}{f(w_n)}\right],
\end{aligned}\right. \tag{18}$$

where ρ and τ are real numbers. Third is (**FS4-1**)

$$\left\{\begin{aligned}
w_n &= x_n + f(x_n),\\
y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]},\\
z_n &= y_n - \frac{f(y_n)}{f[x_n,y_n]+f[y_n,w_n]-f[x_n,w_n]},\\
x_n &= z_n - \frac{f(z_n)}{f[w_n,z_n]}\left[1+\frac{f(z_n)}{f(y_n)}+\frac{f(y_n)}{f(x_n)}+\Big\{2+f[x_n,w_n](3+f[x_n,w_n])\Big\}\left(\frac{f(y_n)}{f(w_n)}\right)^2+\omega\frac{f(z_n)}{f(x_n)}+\phi\frac{f(z_n)}{f(w_n)}\right],
\end{aligned}\right. \tag{19}$$

where ω and ϕ are real numbers. Fourth is (**FS4-2**)

$$\left\{\begin{aligned}
w_n &= x_n - f(x_n),\\
y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]},\\
z_n &= y_n - \frac{f(y_n)}{f[x_n,y_n]+f[y_n,w_n]-f[x_n,w_n]},\\
x_n &= z_n - \frac{f(z_n)}{f[w_n,z_n]}\left[1+\frac{f(z_n)}{f(y_n)}+\frac{f(y_n)}{f(x_n)}+\Big\{2+f[x_n,w_n](-3+f[x_n,w_n])\Big\}\left(\frac{f(y_n)}{f(w_n)}\right)^2\right].
\end{aligned}\right. \tag{20}$$

2. A More general formulation of sixth-order iterative methods

In this section, we go beyond the iterative method (**SK1**) and present a more general formulation which covers most of the sixth-order iterative methods based on one derivative evaluation $f'(x)$. The proposed iterative scheme is (**FD1**)

$$\begin{cases} y_n & = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n & = y_n - G(t_1)\frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} & = z_n - H(t_1,t_2,t_3)\frac{f(z_n)}{f'(x_n)}, \end{cases} \tag{21}$$

where $t_1 = \frac{f(y_n)}{f(x_n)}$, $t_2 = \frac{f(z_n)}{f(x_n)}$ and $t_3 = \frac{f(z_n)}{f(y_n)}$. we state the following theorem.

Theorem 1. Let $f : D \subseteq \Re \to \Re$ be a sufficiently differentiable function, and $\alpha \in D$ is a simple root of $f(x) = 0$, for an open interval D. If x_0 is chosen sufficiently close to α, then iterative scheme (21) converges to α. If G and H satisfy

$$G(0) = 1,\ dG/dt_1(0) = 2,\ H(0,0,0) = 1,\ \partial H/\partial t_1(0,0,0) = 2, \tag{22}$$

and both G, H have bounded higher order derivatives then the iterative scheme (21) has convergence order at-least six and along with conditions (22) if

$$\partial^2 H/\partial t_1^2(0,0,0) = d^2G/dt_1^2(0) + 2,\ \partial H/\partial t_3(0,0,0) = 1 \tag{23}$$

then the iterative scheme (21) has convergence order at least seven.

Proof. let α be the simple root of $f(x) = 0$ and $e_n = x_n - \alpha$. Further we denote $c_1 = f'(\alpha)$ and $c_k = \frac{e_n^k}{k!}\frac{f^{(k)}(\alpha)}{f'(\alpha)}$, where $k = 2, 3, \cdots$. The Tyalor's expression of f and f' around α in terms of e_n:

$$f(x_n) = c_1[e_n + c_2e_n^2 + c_3e_n^3 + c_4e_n^4 + c_5e_n^5 + c_6e_n^6 + O(e_n^7)] \tag{24}$$

$$f'(x_n) = c_1[1 + 2c_2e_n + 3c_3e_n^2 + 4c_4e_n^3 + 5c_5e_n^4 + 6c_6e_n^5 + O(e_n^6)] \tag{25}$$

Substituting (24) and (25) into y_n, we obtain

$$\begin{aligned} y_n = x_n - \frac{f(x_n)}{f'(x_n)},\ y_n - \alpha &= x_n - \alpha - \frac{f(x_n)}{f'(x_n)} = e_n - \frac{f(x_n)}{f'(x_n)} \\ &= c_2e_n^2 + (2c_3 - 2c_2^2)e_n^3 + (3c_4 - 7c_2c_3 + 4c_2^3)e_n^4 + (4c_5 - 10c_2c_4 - 6c_3^2 + 20c_3c_2^2 - 8c_2^4)e_n^5 \\ &+ (-17c_3c_4 + 33c_2c_3^2 - 52c_3c_2^3 + 28c_4c_2^2 - 13c_2c_5 + 5c_6 + 16c_2^5)e_n^6 + O(e_n^7) \end{aligned} \tag{26}$$

Expansion of $f(y_n)$ around α gives

$$\begin{aligned} f(y_n) = c_1[&c_2e_n^2 - 2(-c_3 + c_2^2)e_n^3 + (3c_4 - 7c_2c_3 + 5c_2^3)e_n^4 - 2(-2c_5 + 5c_2c_4 + 3c_3^2 - 12c_3c_2^2 + 6c_2^4)e_n^5 \\ &c_1(37c_2c_3^2 - 73c_3c_2^3 + 28c_2^5 + 34c_4c_2^2 - 17c_3c_4 - 13c_2c_5 + 5c_6)e_n^6 + O(e_n^7)] \end{aligned} \tag{27}$$

We define $G(t_1) = 1 + 2t_1 + M_0t_1^2 + M_1t_1^3 + M_2t_1^4 +$ higher powers in t_1 , where M_0, M_1 and M_2 are finite real numbers. By using (24)-(27), the expressions for z_n and $f(z_n)$ are

$$\begin{aligned} z_n &= y_n - G(t_1)\frac{f(y_n)}{f'(x_n)} \\ z_n - \alpha &= -c_2(-5c_2^2 + M_0c_2^2 + c_3)e_n^4 + (-2c_2c_4 - 2c_3^2 + 32c_3c_2^2 - 36c_2^4 - M_1c_2^4 - 6M_0c_2^2c_3 + 10M_0c_2^4)e_n^5 \\ &(-12M_0c_2c_3^2 - 9M_0c_2^2c_4 + 74M_0c_2^3c_3 - 62M_0c_2^5 - 7c_3c_4 + 66c_2c_3^2 - 262c_3c_2^3 + 48c_4c_2^2 - 3c_2c_5 \\ &+ 170c_2^5 - M_2c_2^5 + 13M_1c_2^5 - 8M_1c_2^3c_3)e_n^6 + O(e_n^7) \end{aligned} \tag{28}$$

$$\begin{aligned} f(z_n) &= -c_1c_2(-5c_2^2 + M_0c_2^2 + c_3)e_n^4 + c_1(-2c_2c_4 - 2c_3^2 + 32c_3c_2^2 - 36c_2^4 - M_1c_2^4 - 6M_0c_2^2c_3 \\ &+ 10M_0c_2^4)e_n^5 + c_1(-12M_0c_2c_3^2 - 9M_0c_2^2c_4 + 74M_0c_2^3c_3 - 62M_0c_2^5 - 7c_3c_4 + 66c_2c_3^2 - 262c_3c_2^3 \\ &+ 48c_4c_2^2 - 3c_2c_5 + 170c_2^5 - M_2c_2^5 + 13M_1c_2^5 - 8M_1c_2^3c_3)e_n^6 + O(e_n^7) \end{aligned} \tag{29}$$

Now, we define $H(t_1, t_2, t_3) = 1 + 2t_1 + R_0t_2 + R_1t_3 + R_2t_1^2 + R_3t_2^2 + R_4t_3^2 +$ higher powers in t_1, t_2, t_3. Again, substituting (25), (28) and (29) in (21), we obtain

$$\begin{aligned} x_{n+1} - \alpha &= z_n - \alpha - H(t_1, t_2, t_3)\frac{f(y_n)}{f'(x_n)} \\ &= -c_2(-5c_2^2 + M_0c_2^2 + c_3)(R_1M_0c_2^2 - 5R_1c_2^2 + 6c_2^2 - R_2c_2^2 + R_1c_3 - c_3)e_n^6 \\ &+ (10R_0c_2^6M_0 - R_0c_2^6M_0^2 + 10R_1M_1c_2^6 - 172R_1c_2^6M_0 + 20R_1M_0^2c_2^6 + 2M_0c_2^3c_4 + 6M_0c_2^2c_3^2 \\ &- 68M_0c_2^4c_3 + 88M_0c_2^6 + 124R_1M_0c_2^4c_3 - 12R_1M_0c_2^2c_3^2 - 10R_1M_0^2c_2^4c_3 - 2R_1M_1c_2^6M_0 - 2R_1M_1c_2^4c_3 \\ &- 2R_0c_2^4M_0c_3 - 4R_1c_4M_0c_2^3 - 22c_4c_2^3 - 66c_3^2c_2^2 + 366c_3c_2^4 + 2c_3^3 - 356c_2^6 + 4c_4c_2c_3 - 6M_1c_2^6 \\ &- 25R_6c_2^6 + 10R_6c_2^6M_0 + 10R_6c_2^4c_3 - R_6c_2^6M_0^2 - R_6c_2^2c_3^2 + M_1c_2^4c_3 + 76R_2c_2^6 - 2R_6c_2^4M_0c_3 \\ &- 4c_2R_1c_4c_3 - 342R_1c_3c_2^4 + 20R_1c_4c_2^3 + 10R_0c_2^4c_3 + 64c_2^2R_1c_3^2 - R_0c_2^2c_3^2 - 25R_0c_2^6 + 360R_1c_2^6 \\ &- 2R_1c_3^3 - 60R_2c_2^4c_3 + 6R_2c_2^2c_3^2 + 2R_2c_2^3c_4 - 18R_2c_2^6M_0 + R_2c_2^6M_1 + 10R_2c_2^4M_0c_3)e_n^7 + O(e_n^8) \end{aligned} \tag{30}$$

By equating one of the factor of coefficient of e_n^6 to zero, we have

$$R_1M_0c_2^2 - 5R_1c_2^2 + 6c_2^2 - R_2c_2^2 + R_1c_3 - c_3 = 0. \tag{31}$$

Now by equating the coefficient of c_3 and c_2^2 in (31), we find

$$\begin{cases} R_1 - 1 = 0, \text{ (coefficient of } c_3) \\ R_1M_0 - 5R_1 + 6 - R_2 = 0, \quad \text{(coefficient of } c_2^2) \end{cases} \tag{32}$$

where $M_0 = 1/2\ d^2G/dt_1^2(0)$, $R_1 = \partial H/\partial t_3(0, 0, 0)$ and $R_2 = 1/2\ \partial^2H/\partial t_1^2(0, 0, 0)$. Which completes the proof. □

Remark 2.1. It is obvious that sixth-order family (21) can construct all the presented iterative methods in (5)-(9) and (11).

(10) can be written as (**FD2**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= x_n - A(t_1)\frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} &= z_n - \frac{f(z_n)}{f[z_n,y_n]+f[z_n,x_n,x_n](z_n-y_n)}, \end{cases} \tag{33}$$

where $t_1 = \frac{f(y_n)}{f(x_n)}$ and $A(t_1) = \left(\frac{1-\theta t_1}{1-t_1}\right)$, with $\theta \in \Re$ and $f[z_n, x_n, x_n] = \frac{f[z_n,x_n]-f'(x_n)}{z_n-x_n}$. The more general form of (33) is stated in the following theorem.

Theorem 2. Let $f : D \subseteq \Re \to \Re$ be a sufficiently differentiable function, and $\alpha \in D$ is a simple root of $f(x) = 0$, for an open interval D. If x_0 is chosen sufficiently close to α, then iterative scheme (33) converges to α. If A satisfies

$$A(0) = 1 \tag{34}$$

then the iterative scheme (33) has convergence order at least six and if along with (34), it satisfies

$$dA/dt_1(0) = 2 \tag{35}$$

then (33) has convergence order at least seven.

Proof. We define $A(t_1) = 1 + M_0t_1 + M_1t_1^2 + M_2t_1^3 +$ higher powers in s. By using the (24)-(27), we obtain

$$\begin{aligned} z_n - \alpha &= y_n - \alpha - (1 + M_0t_1 + M_1t_1^2 + M_2t_1^3)\frac{f(y_n)}{f'(x_n)} \\ &= (-c_2^2(-2 + M_0))e_n^3 + c_2(-9c_2^2 + 7M_0c_2^2 - M_1c_2^2 + 7c_3 - 4M_0c_3)e_n^4 + (10c_2c_4 + 6c_3^2 - 44c_3c_2^2 \\ &+ 30c_2^4 - 6M_0c_2c_4 + 38M_0c_2^2c_3 - 33M_0c_2^4 - M_2c_2^4 - 6M_1c_2^2c_3 + 10M_1c_2^4 - 4M_0c_3^2)e_n^5 + (17c_3c_4 \\ &- 70c_2c_3^2 + 188c_3c_2^3 - 62c_4c_2^2 + 13c_2c_5 - 88c_2^5 + 74M_1c_2^3c_3 - 9M_1c_2^2c_4 - 12M_1c_2c_3^2 - 62M_1c_2^5 \\ &+ 13M_2c_2^5 - 8M_2c_2^3c_3 + 129M_0c_2^5 - 12M_0c_3c_4 + 68M_0c_2c_3^2 - 225M_0c_3c_2^3 + 55M_0c_4c_2^2 - 8M_0c_2c_5)e_n^6 \\ &+ O(e_n^7), \end{aligned} \tag{36}$$

$$\begin{aligned} f(z_n) &= c_1[(-c_2^2(-2 + M_0))e_n^3 + c_2(-9c_2^2 + 7M_0c_2^2 - M_1c_2^2 + 7c_3 - 4M_0c_3)e_n^4 + (10c_2c_4 + 6c_3^2 - 44c_3c_2^2 \\ &+ 30c_2^4 - 6M_0c_2c_4 + 38M_0c_2^2c_3 - 33M_0c_2^4 - M_2c_2^4 - 6M_1c_2^2c_3 + 10M_1c_2^4 - 4M_0c_3^2)e_n^5 + (17c_3c_4 \\ &- 70c_2c_3^2 + 188c_3c_2^3 - 62c_4c_2^2 + 13c_2c_5 - 88c_2^5 + 74M_1c_2^3c_3 - 9M_1c_2^2c_4 - 12M_1c_2c_3^2 - 62M_1c_2^5 \\ &+ 13M_2c_2^5 - 8M_2c_2^3c_3 + 129M_0c_2^5 - 12M_0c_3c_4 + 68M_0c_2c_3^2 - 225M_0c_3c_2^3 + 55M_0c_4c_2^2 - 8M_0c_2c_5)e_n^6 \\ &+ O(e_n^7)]. \end{aligned} \tag{37}$$

By substituting values from (24)-(27), (36) and (37) in x_{n+1} from (33), we get

$$\begin{aligned} x_{n+1}-\alpha &= z_n-\alpha-\frac{f(z_n)}{f[z_n,y_n]+f[z_n,x_n,x_n](z_n-y_n)}\\ e_{n+1} &= c_2^3(-2+M_0)(M_0c_2^2+2c_3-2c_2^2)e_n^6-c_2^2(36c_2^4-46M_0c_2^4+4M_1c_2^4-2M_1c_2^4M_0+14M_0^2c_2^4\\ &\quad -10M_0^2c_3c_2^2-2M_1c_2^2c_3-64c_3c_2^2+57M_0c_2^2c_3+6c_2c_4-3M_0c_2c_4+22c_3^2-12M_0c_3^2)e_n^7+O(e_n^8). \end{aligned} \tag{38}$$

Clearly, if $dA(t_1)/dt_1(0)=M_0=2$ then convergence order is seven otherwise is six. ☐

Now we present derivative-free optimal fourth-order iterative schemes (**FD3**):

$$\begin{cases} w_n &= x_n-\kappa f(x_n),\\ y_n &= x_n-\frac{f(x_n)}{f[x_n,w_n]},\\ x_{n+1} &= y_n-\Big[\frac{g_0}{f[y_n,w_n]}G_0(t_1)+\frac{g_1}{f[y_n,x_n]}G_1(t_2)+\frac{g_2}{f[x_n,w_n]}G_2(t_1,t_2)\Big], \end{cases} \tag{39}$$

where $t_1=\frac{f(y_n)}{f(x_n)}$, $t_2=\frac{f(y_n)}{f(w_n)}$. The convergence of (39) is given in the following theorem.

Theorem 3. Let $f:D\subseteq\Re\to\Re$ be a sufficiently differentiable function, and $\alpha\in D$ is a simple root of $f(x)=0$, for an open interval D. If x_0 is chosen sufficiently close to α, then iterative scheme (39) converges to α. If $G_0(t_1)$, $G_1(t_2)$, $G_2(t_1,t_2)$, g_0, g_1 and g_2 satisfy

$$\begin{cases} G_0(0))=1,\ G_1(0)=1,\ G_2(0,0)=1,\\ dG_0/dt_1(0)=1,\ dG_1/dt_2(0)=1,\ \partial G_2/\partial t_1(0,0)=1,\ \partial G/\partial t_2(0,0)=1,\\ g_0+g_1+g_2=1 \end{cases} \tag{40}$$

and all higher order derivatives of $G_i,\ i=0,1,2$ are bounded then the iterative scheme (39) has convergence order at least four.

Proof. Error term at nth-step is denoted by $e_n=x_n-\alpha$. we call $c_1=f'(\alpha)$ and $c_k=\frac{e_n^k}{k!}\frac{f^{(k)}(\alpha)}{f'(\alpha)}$, where $k=2,3,\cdots$. Taylor's expressions give:

$$f(x_n)=c_1[e_n+c_2e_n^2+c_3e_n^3+c_4e_n^4+c_5e_n^5+c_6e_n^6+O(e_n^7)]. \tag{41}$$

$$\begin{aligned} f(w_n) &= -c_1(-1+\kappa c_1)e_n+c_1c_2(-3\kappa c_1+1+\kappa^2c_1^2)e_n^2-c_1(4\kappa c_1c_3-c_3-3c_3\kappa^2c_1^2+c_3\kappa^3c_1^3+2c_2^2\kappa c_1\\ &\quad -2\kappa^2c_1^2c_2^2)e_n^3+c_1(-5\kappa c_1c_4-5c_2\kappa c_1c_3+8\kappa^2c_1^2c_2c_3+\kappa^2c_1^2c_2^3+c_4+6c_4\kappa^2c_1^2-4c_4\kappa^3c_1^3+c_4\kappa^4c_1^4\\ &\quad -3c_2\kappa^3c_1^3c_3)e_n^4+\cdots+O(e_n^7). \end{aligned} \tag{42}$$

By using (41) and (42), we get

$$\begin{aligned} f[x_n,w_n] &= \frac{f(x_n)-f(w_n)}{x_n-w_n}=-c_1c_2(-2+\kappa c_1)e_n+c_1(3c_3-3\kappa c_1c_3+c_3\kappa^2c_1^2-c_2^2\kappa c_1)e_n^2-c_1(-4c_4\\ &\quad +4c_2\kappa c_1c_3+6\kappa c_1c_4-4c_4\kappa^2c_1^2+c_4\kappa^3c_1^3-2\kappa^2c_1^2c_2c_3)e_n^3+c_1(2\kappa^2c_1^2c_3^2+8\kappa^2c_1^2c_4c_2\\ &\quad -10\kappa c_1c_5+5c_5-7c_4\kappa c_1c_2-3\kappa c_1c_3^2+10c_5\kappa^2c_1^2-5c_5\kappa^3c_1^3+c_5\kappa^4c_1^4-3c_4\kappa^3c_1^3c_2+\kappa^2c_1^2c_2^2c_3)e_n^4\\ &\quad +\cdots+O(e_n^7) \end{aligned} \tag{43}$$

By substituting (41) and (43) in $y=x-f(x_n)/f[x_n,w_n]$, we obtain

$$\begin{aligned} y_n-\alpha &= -c_2(-1+\kappa c_1)e_n^2+(2c_3-3\kappa c_1c_3+c_3\kappa^2c_1^2+2c_2^2\kappa c_1-2c_2^2-\kappa^2c_1^2c_2^2)e_n^3+(3c_4+10c_2\kappa c_1c_3-6\kappa c_1c_4\\ &\quad +4c_4\kappa^2c_1^2-c_4\kappa^3c_1^3-7\kappa^2c_1^2c_2c_3-7c_2c_3-5\kappa c_1c_2^3+2c_2\kappa^3c_1^3c_3+3\kappa^2c_1^2c_2^3+4c_2^3-\kappa^3c_1^3c_2^3)e_n^4+\cdots+O(e_n^7) \end{aligned} \tag{44}$$

Taylor's Expansion of f around y_n by using (44) is

$$\begin{aligned} f(y_n) &= c_1(-c_2(-1+\kappa c_1)e_n^2+(2c_3-3\kappa c_1c_3+c_3\kappa^2c_1^2+2c_2^2\kappa c_1-2c_2^2-\kappa^2c_1^2c_2^2)e_n^3+(3c_4+10c_2\kappa c_1c_3-6\kappa c_1c_4\\ &\quad +4c_4\kappa^2c_1^2-c_4\kappa^3c_1^3-7\kappa^2c_1^2c_2c_3-7c_2c_3-5\kappa c_1c_2^3+2c_2\kappa^3c_1^3c_3+3\kappa^2c_1^2c_2^3+4c_2^3-\kappa^3c_1^3c_2^3)e_n^4+\cdots+O(e_n^7)) \end{aligned} \tag{45}$$

Now, we define

$$\begin{cases} G_0(t_1) &= 1+t_1+L_1t_1^2+\text{higher powers in } t_1,\\ G_1(t_2) &= 1+t_2+M_1t_2^2+\text{higher powers in } t_2,\\ G_2(t_1,t_2) &= 1+t_1+t_2+N_1t_1^2+N_2t_2^2+\text{higher powers in } t_1,\ t_2, \end{cases} \tag{46}$$

where L_1, M_1, N_1 and N_2 are real numbers. By substituting (41)-(45) in $x_{n+1} = y_n - \Big[\frac{g_0}{f[y_n,w_n]}G_0(t_1) + \frac{g_1}{f[y_n,x_n]}G_1(t_2) + \frac{g_2}{f[x_n,w_n]}G_2(t_1,t_2)\Big]$, we obtain

$$\begin{aligned}
e_{n+1} = {} & (-1+g_0+g_1+g_2)c_2(-1+\kappa c_1)e_n^2 + \\
& (-1+g_0+g_1+g_2)(-2c_3+3\kappa c_1c_3 - c_3\kappa^2c_1^2 - 2c_2^2\kappa c_1 + 2c_2^2 + \kappa^2c_1^2c_2^2)e_n^3 + \\
& (-3g_0c_4 - g_0c_2^3 + 6g_0c_2c_3 - 7\kappa^2c_1^2c_2c_3 - 6\kappa c_1c_4 + 3c_4 + 3g_2N_1c_2^3\kappa c_1 - 3g_2N_1c_2^3\kappa^2c_1^2 \\
& + g_2N_1c_2^3\kappa^3c_1^3 + g_2N_2c_2^3\kappa c_1 + g_1M_1c_2^3\kappa c_1 + g_0\kappa^3c_1^3L_1c_2^3 + 3g_0L_1c_2^3\kappa c_1 - 3g_0L_1c_2^3\kappa^2c_1^2 + 3\kappa^2c_1^2c_2^3 \\
& + 4c_4\kappa^2c_1^2 - c_4\kappa^3c_1^3 - 5\kappa c_1c_2^3 - \kappa^3c_1^3c_2^3 + 10c_2\kappa c_1c_3 + 2c_2\kappa^3c_1^3c_3 - 7c_2c_3 + 4c_2^3 - 5g_2\kappa c_1c_2^3 + g_1\kappa^3c_1^3c_2^3 \\
& + 6g_0\kappa c_1c_4 - 4g_0c_4\kappa^2c_1^2 + g_0c_4\kappa^3c_1^3 - 2g_0\kappa c_1c_2^3 + 2g_0\kappa^2c_1^2c_2^3 - g_2N_2c_2^3 - g_1M_1c_2^3 - g_2N_1c_2^3 - g_0L_1c_2^3 \\
& - 8g_0c_2\kappa c_1c_3 + 6g_0\kappa^2c_1^2c_2c_3 - 2g_0c_2\kappa^3c_1^3c_3 + 6g_1\kappa c_1c_4 + 6g_1\kappa^2c_1^2c_2c_3 - 8g_1c_2\kappa c_1c_3 - 2g_1c_2\kappa^3c_1^3c_3 \\
& + 6g_2\kappa^2c_1^2c_2c_3 - 8g_2c_2\kappa c_1c_3 - 2g_2c_2\kappa^3c_1^3c_3 + 6g_1c_2c_3 + 6g_2c_2c_3 - 3g_1c_4 - g_1c_2^3 - 3g_2c_4 + g_2c_2^3 \\
& - g_1\kappa^2c_1^2c_2^3 - 4g_1c_4\kappa^2c_1^2 + g_1c_4\kappa^3c_1^3 + 6g_2\kappa c_1c_4 + 3g_2\kappa^2c_1^2c_2^3 - 4g_2c_4\kappa^2c_1^2 + g_2c_4\kappa^3c_1^3)e_n^4 + \cdots + O(e_n^7)
\end{aligned} \tag{47}$$

Clearly, if $g_0+g_1+g_2=1$ then the coefficients of e_n^2 and e_n^3 are zero and hence the convergence order is four which is optimal in the sense of Kung and Traub. ☐

Remark 2.2. In (39), if we equate $g_0=0$, $g_1=0$, $g_2=1$ and define $G_2(t_1,t_2) = 1+t_1+t_2+\alpha t_1^2+\beta t_2^2$, then the resulted iterative scheme is the first two steps of iterative method (12). Similarly if we equate $g_0=0$, $g_2=0$, $g_1=1$, $\kappa=-1$ and define $G_1(t_2) = (1-\lambda^{-1}t_2)^{-\lambda}$, then it reproduces the first three step of iterative schemes (13) and (14).

In order to extend the iterative scheme (39), we add one more step and get (**FD4**)

$$\begin{cases}
w_n &= x_n - \kappa f(x_n), \\
y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]}, \\
z_n &= y_n - \Big[\frac{(1-g_1-g_2)}{f[y_n,w_n]}G_0(t_1) + \frac{g_1}{f[y_n,x_n]}G_1(t_2) + \frac{g_2}{f[x_n,w_n]}G_2(t_1,t_2)\Big], \\
x_{n+1} &= z_n - \Big[\frac{(1-h_1-h_2-h_3-h_4-h_5)}{f[y_n,z_n]}S_0(t_3,t_4,t_5) + \frac{h_1}{f[z_n,w_n]}S_1(t_1,t_3,t_4,t_5) + \frac{h_2}{f[z_n,x_n]}S_2(t_2,t_3,t_4,t_5) \\
& \quad + \frac{h_3}{f[x_n,y_n]}S_3(t_2,t_3,t_4,t_5) + \frac{h_4}{f[y_n,w_n]}S_4(t_1,t_3,t_4,t_5) + \frac{h_5}{f[x_n,w_n]}S_5(t_1,t_2,t_3,t_4,t_5)\Big]f(z_n),
\end{cases} \tag{48}$$

where $t_1 = \frac{f(y_n)}{f(x_n)}$, $t_2 = \frac{f(y_n)}{f(w_n)}$, $t_3 = \frac{f(z_n)}{f(x_n)}$, $t_4 = \frac{f(z_n)}{f(w_n)}$, $t_5 = \frac{f(z_n)}{f(y_n)}$ and h_1, h_2, h_3, h_4, h_5, $\kappa \neq 0$ are real numbers. The convergence of (48) is discussed in the following theorem.

Theorem 4. By including the statement of (Theorem 3), iterative scheme (48) has convergence order at least six if

$$\begin{cases}
S_0(0,0,0)=1,\ S_1(0,0,0,0)=1,\ S_2(0,0,0,0)=1,\ S_3(0,0,0,0)=1,\ S_4(0,0,0,0)=1, \\
S_5(0,0,0,0,0)=1,\ \partial S_1/\partial t_1(0,0,0,0)=1,\ \partial S_2/\partial t_2(0,0,0,0)=1, \\
\partial S_3/\partial t_2(0,0,0,0)=1,\ \partial S_4/\partial t_1(0,0,0,0)=1, \\
\partial S_5/\partial t_1(0,0,0,0,0)=1,\ \partial S_5/\partial t_2(0,0,0,0,0)=1
\end{cases} \tag{49}$$

and all higher order derivatives of all functions are bounded.

Proof. Taylor's expansion of various orders for S_i, $i=0,1,\cdots,5$ is given below.

$$\begin{cases}
S_0(t_3,t_4,t_5) &= 1+a_1t_3+a_2t_4+a_3t_5 + \text{higher powers in } t_3,\ t_4,\ t_5, \\
S_1(t_1,t_3,t_4,t_5) &= 1+t_1+b_1t_1^2+b_2t_3+b_3t_4+b_4t_5 + \text{higher powers in } t_1,\ t_3,\ t_4,\ t_5, \\
S_2(t_2,t_3,t_4,t_5) &= 1+t_2+b_5t_2^2+b_6t_3+b_7t_4+b_8t_5 + \text{higher powers in } t_2,\ t_3,\ t_4,\ t_5, \\
S_3(t_2,t_3,t_4,t_5) &= 1+t_2+b_9t_2^2+b_{10}t_3+b_{11}t_4+b_{12}t_5 + \text{higher powers in } t_2,\ t_3,\ t_4,\ t_5, \\
S_4(t_1,t_3,t_4,t_5) &= 1+t_1+b_{13}t_1^2+b_{14}t_3+b_{15}t_4+b_{16}t_5 + \text{higher powers in } t_1,\ t_3,\ t_4,\ t_5, \\
S_5(t_1,t_2,t_3,t_4,t_5) &= 1+t_1+t_2+b_{17}t_1^2+b_{18}t_2^2+b_{19}t_3+b_{20}t_4+b_{21}t_5 + \text{higher powers in } t_1,\ t_2,\ t_3,\ t_4,\ t_5,
\end{cases} \tag{50}$$

where a_1, a_2, a_3, b_1, b_2, b_3, b_4, b_5, b_6, b_7, b_8, b_9, b_{10}, b_{11}, b_{12}, b_{13}, b_{14}, b_{15}, b_{16}, b_{17}, b_{18}, b_{19}, b_{20}, and b_{21} are real numbers. By using calculations from (41)-(47) for $x_{n+1} = z_n - \Big[\frac{(1-h_1-h_2-h_3-h_4-h_5)}{f[y_n,z_n]}S_0(t_3,t_4,t_5) + \frac{h_1}{f[z_n,w_n]}S_1(t_1,t_3,t_4,t_5)$
$+ \frac{h_2}{f[z_n,x_n]}S_2(t_2,t_3,t_4,t_5) + \frac{h_3}{f[x_n,y_n]}S_3(t_2,t_3,t_4,t_5) + \frac{h_4}{f[y_n,w_n]}S_4(t_1,t_3,t_4,t_5) + \frac{h_5}{f[x_n,w_n]}S_5(t_1,t_2,t_3,t_4,t_5)\Big]f(z_n)$,

we obtain the following error equation.

$$\begin{aligned}e_{n+1} &= -(2h_4b_{13}c_2^2\kappa c_1 - h_4b_{13}c_2^2\kappa^2c_1^2 + a_3g_1M_1c_2^2 - h_5b_{18}c_2^2 - h_5b_{17}c_2^2 - h_3b_9c_2^2 + h_5b_{21}g_1\kappa^2c_1^2c_2^2\\
&+ h_5b_{21}g_2c_2^2\kappa c_1 - 2h_5b_{21}L_1c_2^2\kappa c_1 + h_5b_{21}L_1c_2^2\kappa^2c_1^2 + 5c_2^2h_5 + h_4\kappa^2c_1^2c_2^2 + h_4\kappa c_1c_3 - 4h_4c_2^2\kappa c_1 + h_1\kappa^2c_1^2c_2^2\\
&+ h_1\kappa c_1c_3 - 3h_1c_2^2\kappa c_1 - c_2^2h_2\kappa c_1 - 2c_2^2h_3\kappa c_1 - 5c_2^2h_5\kappa c_1 + h_2\kappa c_1c_3 + h_3\kappa c_1c_3 + h_5\kappa^2c_1^2c_2^2 + h_5\kappa c_1c_3\\
&- a_3\kappa^2c_1^2c_2^2 - a_3\kappa c_1c_3 + 4a_3c_2^2\kappa c_1 + 2a_3h_1g_2c_2^2 + \cdots + h_3b_{12}L_1c_2^2\kappa^2c_1^2 - 2h_2b_8g_1c_2^2\kappa c_1 + h_2b_8g_1\kappa^2c_1^2c_2^2\\
&+ h_2b_8g_2c_2^2\kappa c_1 - 2h_2b_8L_1c_2^2\kappa c_1 + h_2b_8L_1c_2^2\kappa^2c_1^2)c_2(-1+\kappa c_1)(-g_1\kappa^2c_1^2c_2^2 - L_1c_2^2\kappa^2c_1^2 - g_2N_1c_2^2\kappa^2c_1^2\\
&+ \kappa^2c_1^2c_2^2 + g_1L_1c_2^2\kappa^2c_1^2 + g_2L_1c_2^2\kappa^2c_1^2 - 4c_2^2\kappa c_1 + 2g_1c_2^2\kappa c_1 + 2g_2N_1c_2^2\kappa c_1 - g_2c_2^2\kappa c_1 - 2g_1L_1c_2^2\kappa c_1\\
&+ \kappa c_1c_3 + 2L_1c_2^2\kappa c_1 - 2g_2L_1c_2^2\kappa c_1 - c_3 - g_2N_2c_2^2 - g_2N_1c_2^2 + 2g_2c_2^2 + 3c_2^2 - g_1M_1c_2^2 - L_1c_2^2 + g_1L_1c_2^2\\
&+ g_2L_1c_2^2)e_n^6 + O(e_n^7)\end{aligned} \tag{51}$$

□

Remark 2.3. In (48), by equating $\kappa = -1$, $g_1 = 1$, $g_2 = 0$, $h_3 = 1$, $h_1 = h_2 = h_4 = h_5 = 0$ and defining $G_2(t_2) = (1-\lambda^{-1}t_2)^{-\lambda}$, $S_3(t_2,t_3,t_4,t_5) = (1-\lambda^{-1}t_2)^{-\lambda}$ gives the iterative scheme (13), by setting $\kappa = -1$, $g_1 = 1$, $g_2 = 0$, $h_1 = h_2 = h_3 = h_4 = h_5 = 0$ and defining $G_2(t_2) = (1-\lambda^{-1}t_2)^{-\lambda}$,$S_0(t_3,t_4,t_5) = 1$, we obtain the iterative schemes (14). (12) can be constructed by choosing parameters $g_1 = 0$, $g_2 = 1$, $h_5 = 1$, $h_1 = h_2 = h_3 = h_4 = 0$ and weight functions $G_2(t_1,t_2) = 1 + t_1 + t_2 + \alpha t_1^2 + \beta t_2^2$, $S_5(t_1,t_2,t_3,t_4,t_5) = 1 + t_1 + t_2 + \alpha t_1^2 + \beta t_2^2 + \eta t_5$.

F. Soleymani constructs derivative-free sixth-order iterative scheme (16). In the following theorem, we present a modified iterative family with convergence order seven.

Theorem 5. Let $f : D \subseteq \Re \to \Re$ be a sufficiently differentiable function, and $\alpha \in D$ is a simple root of $f(x) = 0$, for an open interval D. If x_0 is chosen sufficiently close to α, then iterative scheme (**FD5**)

$$\begin{cases} w_n &= x_n - \kappa f(x_n),\\ y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]},\\ z_n &= y_n - \left[\frac{1-g_1-g_2}{f[y_n,w_n]}G_0(t_1) + \frac{g_1}{f[x_n,y_n]}G_1(t_2) + \frac{g_2}{f[x_n,w_n]}G_2(t_1,t_2)\right]f(y_n),\\ x_{n+1} &= z_n - H(t_3,t_4,t_5)\frac{f(z_n)}{f[z_n,y_n]-(h-1)f[z_n,w_n]+hf[z_n,x_n]-hf[y_n,x_n]+(h-1)f[y_n,w_n]},\end{cases} \tag{52}$$

where

$$\begin{cases} G_0(t_1) &= 1 + t_1 + a_1t_1^2 + \text{higher powers in } t_1,\\ G_1(t_2) &= 1 + t_2 + a_2t_2^2 + \text{higher powers in } t_2,\\ G_2(t_1,t_2) &= 1 + t_1 + t_2 + a_3t_1^2 + a_4t_2^2 + a_5t_1t_2 + \text{higher powers in } t_1,\ t_2,\\ H(t_3,t_4,t_5) &= 1 + a_6t_3 + a_7t_4 + a_8t_5 + \text{higher powers in } t_3,\ t_4,\ t_5,\end{cases} \tag{53}$$

t_i, $i = 1,2,..,5$ are consistent with previous definition in (48), g_1, g_2, h, a_j, $j = 1,\cdots,8$ are real numbers.

has convergence order at-least six if

$$\begin{cases} G_0(0) = 1,\ G_1(0) = 1,\ G_2(0,0) = 1,\ H(0,0,0) = 1,\\ dG_0/dt_1(0) = 1,\ dG_1/dt_2(0) = 1,\ \partial G_2/\partial t_1(0,0) = 1,\ \partial G_2/\partial t_2(0,0) = 1,\end{cases} \tag{54}$$

and has convergence order seven if along with (54), the following condition is valid

$$\partial H/\partial t_5(0,0,0) = 0. \tag{55}$$

Proof. The terminology of (Theorem 3) and expressions in (41), (42) assist to perform all necessary calculations by using Maple 13 [23] software-package in order to get the following error equation:

$$\begin{aligned}e_{n+1} &= \boldsymbol{a_8}c_2(-1+\kappa c_1)(\kappa^2c_1^2c_2^2 - c_3 + \kappa c_1c_3 + 3c_2^2 - 4c_2^2\kappa c_1 + g_1a_1c_2^2 - a_1c_2^2 + 2a_1c_2^2\kappa c_1 - a_1c_2^2\kappa^2c_1^2 - g_2c_2^2\kappa c_1\\
&+ 2g_1c_2^2\kappa c_1 - g_1\kappa^2c_1^2c_2^2 + g_2a_1c_2^2 - g_1a_2c_2^2 - g_2a_3c_2^2 - g_2a_5c_2^2 - g_2a_4c_2^2 + 2g_2c_2^2 - 2g_1a_1c_2^2\kappa c_1 + g_1a_1c_2^2\kappa^2c_1^2\\
&- 2g_2a_1c_2^2\kappa c_1 + g_2a_1c_2^2\kappa^2c_1^2 + 2g_2a_3c_2^2\kappa c_1 - g_2a_3c_2^2\kappa^2c_1^2 + g_2a_5c_2^2\kappa c_1)^2\boldsymbol{e_n^6}\\
&+ -(\kappa^2c_1^2c_2^2 - c_3 + \kappa c_1c_3 + 3c_2^2 - 4c_2^2\kappa c_1 + g_1a_1c_2^2 - a_1c_2^2 + 2a_1c_2^2\kappa c_1 - a_1c_2^2\kappa^2c_1^2 - g_2c_2^2\kappa c_1 + 2g_1c_2^2\kappa c_1\\
&- g_1\kappa^2c_1^2c_2^2 + g_2a_1c_2^2 - g_1a_2c_2^2 - g_2a_3c_2^2 - g_2a_5c_2^2 - g_2a_4c_2^2 + 2g_2c_2^2 - 2g_1a_1c_2^2\kappa c_1 + g_1a_1c_2^2\kappa^2c_1^2\\
&+ \cdots +\\
&- a_6a_1c_2^4 - 2a_8c_3^2 - 30a_8c_2^4 + 3a_6c_2^4 - 3a_6\kappa^2c_1^2c_2^2c_3 + a_6c_3\kappa^3c_1^3c_2^2 + 3a_6\kappa c_1c_2^2c_3 - 5a_6g_1\kappa^2c_1^2c_2^4\\
&+ 2a_6g_1\kappa c_1c_2^4 + 4a_6g_1\kappa^3c_1^3c_2^4 + 4a_6g_2\kappa^2c_1^2c_2^4 - 5a_6g_2\kappa c_1c_2^4 - a_6g_2\kappa^3c_1^3c_2^4 - a_6g_1\kappa^4c_1^4c_2^4 - a_6a_1c_2^4\kappa^4c_1^4\\
&+ 4a_6a_1c_2^4\kappa^3c_1^3)\boldsymbol{e_n^7} + O(e_n^8)\end{aligned} \tag{56}$$

Clearly, $a_8 = \partial H/\partial t_5(0,0,0)$ is a factor in asymptotic error constant of e_n^6, if $a_8 = 0$ then (53) has convergence order seven otherwise convergence order is six. □

Remark 2.4. Iterative scheme (53) reproduces F. Soleymani derivative-free sixth-order iterative scheme (16) if

$$h = 0,\ g_1 = g_2 = 0,\ G_0(t_1) = 1,\ H(t_3,t_4,t_5) = 1. \tag{57}$$

The derivative-free version of (33) is (**FD6**):

$$\begin{cases} w_n &= x_n - \kappa f(x_n), \\ y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]}, \\ z_n &= y_n - \left[\frac{1-g_1-g_2}{f[y,w]}G_0(t_1) + \frac{g_1}{f[x_n,y_n]}G_1(t_2) + \frac{g_2}{f[x_n,w_n]}G_2(t_1,t_2)\right], \\ x_{n+1} &= z_n - H(t_3,t_4,t_5)\frac{f(z_n)}{f[z_n,y_n]+(f[z_n,x_n]-f[w_n,x_n])\frac{z_n-y_n}{z_n-x_n}}, \end{cases} \tag{58}$$

where

$$\begin{cases} G_0(t_1) &= 1 + t_1 + a_1 t_1^2 + \text{higher powers in } t_1, \\ G_1(t_2) &= 1 + t_2 + a_2 t_2^2 + \text{higher powers in } t_2, \\ G_2(t_1,t_2) &= 1 + t_1 + t_2 + a_3 t_1^2 + a_4 t_2^2 + a_5 t_1 t_2 + \text{higher powers in } t_1,\ t_2, \\ H(t_3,t_4,t_5) &= 1 + a_6 t_3 + a_7 t_4 + a_8 t_5 + \text{higher powers in } t_3,\ t_4,\ t_5, \end{cases} \tag{59}$$

t_i, $i = 1,2,\cdots,5$ are consistent with previous definition in (48), g_1, g_2, h, a_j, $j = 1,\cdots,8$ are real numbers.

Theorem 6. Let $f : D \subseteq \Re \to \Re$ be a sufficiently differentiable function, and $\alpha \in D$ is a simple root of $f(x) = 0$, for an open interval D. If x_0 is chosen sufficiently close to α, then iterative scheme (59) has convergence order at-least six if

$$\begin{cases} G_0(0) = 1,\ G_1(0) = 1,\ G_2(0,0) = 1,\ H(0,0,0) = 1, \\ dG_0/dt_1(0) = 1,\ dG_1/dt_2(0) = 1,\ \partial G_2/\partial t_1(0,0) = 1,\ \partial G_2/\partial t_2(0,0) = 1. \end{cases} \tag{60}$$

Proof. We obtain the following error equation

$$\begin{aligned} e_{n+1} = {} & (-1+\kappa c_1)(\kappa^2 c_1^2 c_2^2 - c_3 + \kappa c_1 c_3 + 3c_2^2 - 4c_2^2\kappa c_1 + g_1 a_1 c_2^2 + g_2 a_1 c_2^2 - g_1 a_2 c_2^2 - g_2 c_2^2 a_5 - g_2 a_4 c_2^2 \\ & - g_2 a_3 c_2^2 - a_1 c_2^2 - 2g_1 a_1 c_2^2 \kappa c_1 + g_1 a_1 c_2^2 \kappa^2 c_1^2 - 2g_2 a_1 c_2^2 \kappa c_1 + g_2 a_1 c_2^2 \kappa^2 c_1^2 + 2g_2 a_3 c_2^2 \kappa c_1 + 2a_1 c_2^2 \kappa c_1 \\ & - a_1 c_2^2 \kappa^2 c_1^2 + 2g_1 c_2^2 \kappa c_1 - g_1 \kappa^2 c_1^2 c_2^2 - g_2 c_2^2 \kappa c_1 + 2g_2 c_2^2 - g_2 a_3 c_2^2 \kappa^2 c_1^2 + g_2 c_2^2 a_5 \kappa c_1)(\kappa^2 c_1^2 c_2^2 - c_2^2 \kappa c_1 + 3c_2^2 a_8 \\ & - a_8 c_3 + 2c_2^2 a_8 g_2 - a_1 c_2^2 a_8 - c_2^2 a_8 a_2 g_1 - c_2^2 a_8 g_2 a_3 - c_2^2 a_8 g_2 a_5 - c_2^2 a_8 g_2 a_4 + c_1 a_8 c_3 \kappa + a_1 c_2^2 a_8 g_1 \\ & + a_1 c_2^2 a_8 g_2 - c_1^2 c_2^2 a_8 \kappa^2 g_1 + 2c_1 c_2^2 a_8 \kappa g_1 - c_1 c_2^2 \kappa a_8 g_2 - c_1^2 c_2^2 \kappa^2 a_8 g_2 a_3 + c_1 c_2^2 \kappa a_8 g_2 a_5 + 2c_1 c_2^2 \kappa a_8 g_2 a_3 \\ & - a_1 c_1^2 c_2^2 a_8 \kappa^2 + 2a_1 c_1 c_2^2 a_8 \kappa + a_1 c_1^2 c_2^2 a_8 \kappa^2 g_1 + a_1 c_1^2 c_2^2 \kappa^2 a_8 g_2 - 2a_1 c_1 c_2^2 a_8 \kappa g_1 - 2a_1 c_1 c_2^2 \kappa a_8 g_2 + c_1^2 c_2^2 a_8 \kappa^2 \\ & - 4c_1 c_2^2 a_8 \kappa)c_2 e_n^6 + O(e_n^7) \end{aligned} \tag{61}$$

□

3. Numerical computations

Definition 1. Let x_{n-1}, x_n and x_{n+1} be successive iterations closer to the root α of $f(x) = 0$, the computational order of convergence (COC) [24], can be approximated by

$$COC \approx \frac{ln|(x_{n+1}-\alpha)(x_n-\alpha)^{-1}|}{ln|(x_n-\alpha)(x_{n-1}-\alpha)^{-1}|}. \tag{62}$$

A set of twelve functions is listed in Table 1 which is taken from [25] to validate the iterative methods and performance. In all methods, twelve total number of function evaluations (TNFE) are used.

3.1. Sixth-order convergence performance evaluation of FD1.

For the purpose of comparison between **FD1** and **SG**, **NT1**, **NT2**, **CH**, **GR**, **AL**, we derive an iterative method from (21) which is given as (**FD1-M1**)

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= y_n - G(t_1)\frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} &= z_n - H_1(t_1)H_2(t_2)H_3(t_3)\frac{f(z_n)}{f'(x_n)}, \end{cases} \tag{63}$$

where $t_1 = \frac{f(y_n)}{f(x_n)}, t_2 = \frac{f(z_n)}{f(x_n)}, t_3 = \frac{f(z_n)}{f(y_n)}, G(t_1) = \frac{1}{1-2t_1}, H_1(t_1) = \frac{1}{1-2t_1-t_1^2}, H_2(t_2) = 1+2t_2$ and $H_3(t_3) = \frac{1}{1-1.1t_3}$. Table 2 shows the performance of different listed iterative methods in terms of absolute error ($|x_n - \alpha|$). Clearly **FD1-M1** is competitive with other developed methods and freedom to choose different weight functions, makes it more accurate and stable in the third step of iterative scheme (21).

TABLE 1. List of test functions

Functions	Roots
$f_1(x) = exp(x)\ sin(x) + ln(1 + x^2)$	$\alpha = 0$
$f_2(x) = x^{15} + x^4 + 4x^2 - 15$	$\alpha = 1.148538...$
$f_3(x) = (x - 2)(x^{10} + x + 1)\ exp(-x - 1)$	$\alpha = 2$
$f_4(x) = exp(-x^2 + x + 2) - cos(x + 1) + x^3 + 1$	$\alpha = -1$
$f_5(x) = (x + 1)\ exp(sin(x)) - x^2\ exp(cos(x)) - 1$	$\alpha = 0$
$f_6(x) = sin(x)^2 - x^2 + 1$	$\alpha = 1.40449165...$
$f_7(x) = 10\ exp(-x^2) - 1$	$\alpha = 1.517427...$
$f_8(x) = (x^2 - 1)^{-1} - 1$	$\alpha = 1.414214...$
$f_9(x) = ln(x^2 + x + 2) - x + 1$	$\alpha = 4.15259074...$
$f_{10}(x) = cos(x)^2 - x/5$	$\alpha = 1.08598268...$
$f_{11}(x) = x^{10} - 2x^3 - x + 1$	$\alpha = 0.591448093...$
$f_{12}(x) = exp(sin(x)) - x + 1$	$\alpha = 2.63066415...$

TABLE 2. Numerical comparison between absolute errors($|x_n - \alpha|, TNFE = 12$)

$f_n(x)$,x_0	FD1-M1	SG	NT1	NT2	CH	GR	AL
		$a = -1$	$a = -1$				$\theta = -1.01$
f_1, 0.25	**2.71 e-142**	2.82e-121	1.22e-127	4.25e-84	5.12e-82	9.23 e-93	6.22e-141
f_2 , 1.1	**3.90 e-190**	3.72 e-153	1.23 e-122	1.47 e-95	2.45 e-104	4.43 e-141	1.60 e-123
f_3, 2.1	**2.31e-181**	3.75e-135	6.27e-135	2.89e-93	1.07e-94	6.52e-109	1.80e-147
f_4, -0.5	**5.28 e-218**	8.50 e-181	5.30 e-143	2.70 e-173	1.27 e-174	5.89 e-207	8.99 e-191
f_5, 0.25	**1.96e-247**	4.05e-184	2.51e-165	3.77e-213	7.33e-162	2.97e-243	2.20e-215
f_6, 1.2	**3.79e-215**	1.41e-181	7.22e-173	3.90e-123	1.69e-123	6.80e-154	2.42e-180
f_7, 1.0	**5.26e-123**	2.88e-95	2.10e-113	3.87e-69	6.28e-67	1.81e-76	2.55e-118
f_8, 1.6	**2.27e-181**	1.18e-157	1.65e-97	6.69e-59	1.23e-103	2.26e-147	4.14e-60
f_9, 4.4	2.92e-402	3.13e-366	4.80e-387	2.35e-344	6.52e-341	1.89e-353	**2.41e-432**
f_{10}, 1.5	2.33e-56	2.22e-53	3.42e-64	3.58e-31	10.1	5.45e-56	**8.42e-58**
f_{11}, 0.25	**8.99e-115**	1.30e-97	1.54e-91	1.31e-53	3.86e-45	4.77e-89	1.72e-92
f_{12}, 2.0	**3.53e-173**	1.80e-132	3.42e-138	1.84e-117	2.78e-115	3.52e-126	8.05e-162

TABLE 3. Computational order of convergence (COC)

$f_n(x)$	FD1-M1	SG	NT1	NT2	CH	GR	AL
f_1	5.9999	5.9994	6.0003	5.9989	5.9987	5.9995	5.9965
f_2	6.0000	6.0000	5.9999	5.9997	5.9998	6.0000	5.9933
f_3	6.0000	6.0000	5.9999	5.9995	5.9996	5.9999	6.0012
f_4	6.0000	6.0000	6.0000	6.0000	6.0000	6.0000	5.9998
f_5	6.0000	6.0000	5.9999	6.0000	5.9999	6.0000	5.9999
f_6	6.0000	6.0000	5.9999	5.9999	5.9999	6.0000	5.9998
f_7	5.9999	5.9992	6.0001	5.9974	5.9972	5.9991	5.9860
f_8	6.0000	6.0000	6.0001	5.9985	5.9999	6.0000	6.0971
f_9	6.0000	6.0000	6.0000	6.0000	6.0000	6.0000	6.0000
f_{10}	5.9806	6.0038	6.0098	5.8343	3.4125	5.9880	6.1207
f_{11}	5.9999	5.9981	6.0001	5.9909	5.9856	6.0000	6.0120
f_{12}	6.0000	6.0000	6.0000	6.0002	6.0002	6.0001	5.9981

3.2. **Seventh-order convergence performance evaluation of FD1.** To compare the results, we construct seventh-order iterative scheme (**FD1-M2**) from (21):

$$\begin{cases} y_n &= x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n &= y_n - G(t_1)\frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} &= z_n - H_1(t_1)H_2(t_2)H_3(t_3)\frac{f(z_n)}{f'(x_n)}, \end{cases} \tag{64}$$

where $t_i's$ are defined previously, $G(t_1) = \frac{1}{1-2t_1}, H_1(t_1) = \frac{1}{1-2t_1-t_1^2}, H_2(t_2) = 1 + 2.1t_2$ and $H_3(t_3) = \frac{1}{1-t_3}$. In (33), if we define $A(t_1) = \frac{1+t_1}{1-t_1}$ then (33) is a seventh-order iterative scheme (**AL1**) which is developed in [17] and if $A(t_1) = \frac{1}{(1-t_1)^2}$ then again (33) is a seventh-order iterative scheme (**FD2-M1**).

Table 4. Numerical comparison between absolute errors($|x_n - \alpha|, TNFE = 12$) and COC

$f_n(x)$,x_0	FD1-M2	FD2-M1	AL1	FD1-M2 (COC)	FD2-M1 (COC)	AL1 (COC)
f_1, 0.25	**3.60e-182**	8.55e-177	6.80e-167	6.9998	6.9999	6.9997
f_2, 1.1	**2.99e-263**	6.71e-172	1.63e-145	7.0000	7.0000	6.9999
f_3, 2.1	**8.60e-223**	3.95e-177	3.11e-178	7.0000	7.0000	7.0000
f_4, -0.5	**2.78e-297**	4.95e-247	1.45e-233	7.0000	7.0000	7.0000
f_5, 0.25	**1.31e-307**	1.45e-306	2.81e-273	7.0000	7.0000	7.0000
f_6, 1.2	**5.02e-304**	2.70e-246	1.27e-222	7.0000	7.0000	7.0000
f_7, 1.0	**1.51e-156**	1.83e-145	3.98e-136	6.9997	6.9998	6.9996
f_8, 1.6	**4.95e-240**	3.58e-102	2.35e-67	6.9995	7.0001	6.9978
f_9, 4.4	**5.10e-662**	3.74e-567	1.07e-561	7.0000	7.0000	7.0000
f_{10}, 1.5	1.08e-77	**1.33e-82**	1.73e-64	6.8846	7.0083	6.9701
f_{11}, 0.25	**8.67e-162**	4.51e-141	1.31e-103	7.0000	7.0003	7.0019
f_{12}, 2.0	**1.30e-242**	4.29e-196	9.45e-193	7.0000	7.0000	6.9999

Table 4 shows absolute error due to method **FD2-M1** has small magnitude in comparison with other describe methods for the set of given twelve test functions and initial guesses. Only in the case of f_9 and f_{10}, **AL** has better absolute error magnitude.

3.3. **Sixth-order convergence performance evaluation of FD4, FD5, FD6.** For comparison, we develop some particular cases of derivative-free methods. In all calculations the value of parameter $\kappa = 1/100$ and $t_1 = \frac{f(y_n)}{f(x_n)}$, $t_2 = \frac{f(y_n)}{f(w_n)}$, $t_3 = \frac{f(z_n)}{f(x_n)}$, $t_4 = \frac{f(z_n)}{f(w_n)}$, $t_5 = \frac{f(z_n)}{f(y_n)}$.

- **FD4**

$$\begin{cases} w_n &= x_n - \kappa f(x_n), \\ y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]}, \\ z_n &= y_n - \frac{1+t_1-t_2}{1-2t_2}\frac{f(y_n)}{f[x_n,w_n]}, \\ x_{n+1} &= z_n - \frac{1}{2}\left[\frac{1}{1-t_5-2t_3-2t_4}\frac{1}{f[z_n,y_n]} + \frac{1-2t_2}{1-3t_2}\frac{1}{f[z_n,x_n]}\right]f(z_n). \end{cases} \tag{65}$$

- **FD5**

$$\begin{cases} w_n &= x_n - \kappa f(x_n), \\ y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]}, \\ z_n &= y_n - \frac{1}{1-t_2}\frac{f(y_n)}{f[x_n,y_n]}, \\ x_{n+1} &= z_n - \left(1 - \frac{t_5}{10}\right)^{-1}\frac{f(z_n)}{f[z_n,y_n]+f[z_n,w_n]-f[y_n,w_n]}. \end{cases} \tag{66}$$

- **FD6**

$$\begin{cases} w_n &= x_n - \kappa f(x_n), \\ y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]}, \\ z_n &= y_n - \frac{1-t_1+t_2}{1-2t_1}\frac{f(y_n)}{f[x_n,w_n]}, \\ x_{n+1} &= z_n - \left(1 - \frac{t_5}{10}\right)^{-1}\frac{f(z_n)}{f[z_n,y_n]+(f[z_n,x_n]-f[x_n,w_n])\frac{z_n-y_n}{z_n-x_n}}. \end{cases} \tag{67}$$

- **SK2M1, SK2M2**
  In [18], Authors developed two methods by selecting different values of parameters. If $\alpha = \beta = \frac{5}{2}$, $\eta = 1$ then **SK2** is **SK2M1** and if $\alpha = \beta = \eta = 1$ then **SK2** is **SK2M2**.

Table 5 shows overall performance of derivative-free sixth-order iterative methods **FD4, FD5, FD6**. Absolute error $|x_n - \alpha|$, for developed sixth-order derivative-free methods, is comparatively better than referenced sixth-order derivative-free iterative methods. Computational order of convergence is given in Table 6. **dgt** stands for **divergent** and **X** is for **no information**.

3.4. **Seventh-order convergence performance evaluation of FD7. FD7** is seventh-order convergent iterative method which is deduced from **FD5** and is defined as

$$\begin{cases} w_n &= x_n - \kappa f(x_n), \\ y_n &= x_n - \frac{f(x_n)}{f[x_n,w_n]}, \\ z_n &= y_n - \frac{1-2t_1}{1-3t_1}\frac{f(y_n)}{f[y_n,w_n]}, \\ x_{n+1} &= z_n - \frac{1}{1-t_3}\frac{f(z_n)}{f[z_n,y_n]+f[z_n,x_n]-f[x_n,y_n]}. \end{cases} \tag{68}$$

Table 5. Numerical comparison between absolute errors($|x_n - \alpha|, TNFE = 12$)

$f_n(x)$	FD4	FD5	FD6	TS1	TS2	SK2M1	SK2M2	FS1	FS2
f_1, 0.25	**4.e-172**	2.e-158	**8.e-172**	9.e-54	2.e-66	3.e-103	4.e-83	1.e-59	2.e-104
f_2, 1.1	1.e-190	2.e-297	6.e-221	0.01	3.e-14	7.e-162	7.e-241	2.e-7	**9.e-333**
f_3, 2.1	2.e-107	**2.e-203**	4.e-168	3.e-18	4.e-24	2.e-197	6.e-145	3.e-7	**5.e-204**
f_4, 0.5	**3.e-220**	2.e-198	8.e-213	3.e-127	2.e-202	1.e-217	3.e-175	9.e-196	4.e-171
f_5, 0.25	**3.e-247**	**7.e-248**	6.e-215	2.e-129	8.e-172	4.e-157	1.e-197	2.e-171	3.e-215
f_6, 1.2	1.e-200	2.e-186	**5.e-247**	1.e-132	5.e-153	8.e-112	2.e-161	8.e-194	2.e-161
f_7, 1.0	3.e-140	8.e-130	**8.e-146**	dgt	dgt	3.e-78	1.e-65	dgt	6.e-87
f_8, 1.6	**3.e-168**	1.e-90	**8.e-165**	5.e-55	2.e-71	2.	5.e-25	2.e-68	3.e-145
f_9, 4.4	**5.e-466**	3.e-392	1.e-404	2.e-395	2.e-424	1.e-383	1.e-366	2.e-429	5.e-378
f_{10}, 1.5	1.e-70	6.e-70	8.e-65	1.e-138	**3.e-168**	0.06	1.	9.e-156	5.e-68
f_{11}, 0.25	7.e-118	4.e-118	**4.e-169**	2.e-71	1.e-105	1.	8.e-29	1.e-118	3.e-85
f_{12}, 2.0	1.e-189	3.e-149	1.e-144	8.e-137	2.e-194	2.e-173	2.e-144	**5.e-215**	4.e-191

Table 6. Computational order of convergence (COC)

$f_n(x)$	FD4	FD5	FD6	TS1	TS2	SK2M1	SK2M2	FS1	FS2
f_1	6.01	6.00	6.0	5.99	6.00	5.98	6.00	5.98	6.00
f_2	6.00	6.00	6.0	-1.15	6.24	6.00	6.00	3.33	6.00
f_3	6.00	6.00	6.0	5.32	5.63	6.00	6.00	3.81	6.00
f_4	6.00	6.00	6.0	6.00	6.00	6.00	6.00	6.00	6.00
f_5	6.00	6.00	6.0	6.00	6.00	6.00	6.00	6.00	6.00
f_6	6.00	6.00	6.0	6.00	6.00	6.00	6.00	6.00	6.00
f_7	6.00	6.00	6.0	X	X	5.98	6.00	X	6.00
f_8	6.01	6.00	6.0	5.99	6.00	5.23	5.78	6.00	6.00
f_9	6.00	6.00	6.0	6.00	6.00	6.00	6.00	6.00	6.00
f_{10}	6.30	6.01	6.0	6.00	6.00	1.51	0.00299	6.00	5.99
f_{11}	6.05	6.00	6.0	6.00	6.00	2.57	5.91	6.00	6.00
f_{12}	5.98	6.00	6.0	6.00	6.00	6.00	6.00	6.00	6.00

Table 7. Numerical comparison between absolute errors($|x_n - \alpha|, TNFE = 12$) and COC

$f_n(x)$	FD7(κ)(COC)	FS3-1(COC)	FS3-2(COC)	FS4-1(COC)	FS4-2(COC)
f_1, 0.25	**2.45e-378**(1.0)(10)	3.23e-99(7)	4.83e-318(9)	2.96e-85(7)	6.60e-311(10)
f_2, 1.1	**1.22e-388**(0.01)(7)	3.33e-31(6.42)	8.14e-24(7.02)	2.37e-7(3.30)	2.04(-0.0211)
f_3, 2.1	**3.29e-266**(0.01)(7)	1.37e-40(6.80)	1.32e-26(7.42)	3.96e-8(4.18)	divergent(X)
f_4, 0.5	**8.61e-262**(0.01)(7)	1.89e-221(7)	1.17e-144(7)	2.06e-210(7)	8.75e-75(6.97)
f_5, 0.25	1.54e-303(0.01)(7)	2.47e-220(7)	**6.43e-397**(7)	1.33e-206(7)	1.25e-384(7)
f_6, 1.2	**5.15e-353**(0.01)(7)	1.71e-251(7)	6.17e-74(7.01)	1.34e-257(7)	4.81e-44(7.05)
f_7, 1.0	**3.65e-213**(0.01)(7)	0.320(4.96)	3.03(X)	dgt(X)	3.03(X)
f_8, 1.6	1.10e-174(0.01)(7)	**1.09e-234**(7)	7.46e-22(7.09)	2.68e-233(7)	290(1.43)
f_9, 4.4	**9.09e-689**(0.01)(7)	4.55e-617(7)	1.71e-517(7)	5.06e-640(7)	4.45e-506(7)
f_{10}, 1.5	**3.81e-262**(-1.0)(7)	6.26e-174(7)	3.87e-5(2.04)	1.96e-214(7)	7.16(X)
f_{11}, 0.25	**7.21e-212**(0.01)(7)	1.33e-125(7)	5.57e-77(6.99)	6.28e-147(7)	1.58e-52(6.97)
f_{12}, 2.0	**3.52e-271**(0.01)(7)	7.93e-257(7)	6.79e6(-7.79)	3.40e-222(7)	2.64(-0.261)

All parameters set to zero value namely $\gamma = 0$, $\delta = 0$ in **FS3-1**, $\rho = 0$, $\tau = 0$ in **FS3-2** and $\omega = 0$, $\phi = 0$ in **FS4**. Table 7 represents the comparison between absolute errors as well as computational order of convergence. Clearly **FD7** is superior than other methods in comparison.

4. CONCLUSIONS

We have presented derivative-based and derivative-free iterative methods by introducing free weighting parameters g_0, g_1, g_2 and weight functions. Weighting parameters, in the second step of derivative-free iterative methods, give flexibility for selecting different forms of approximation for first order derivative. A combination of derivative approximations can be constructed by fixing weighting parameters. The proper selection weight functions for both derivative-free and derivative-based iterative methods help to improve the accuracy. The presented iterative methods are defined in very general form which actually reproduce many existing iterative methods. By introducing different form of weighting functions and selection of parameters produce new families of iterative methods. We explore some of them to show the effectiveness of newly constructed iterative methods.

Acknowledgement. This research was supported by Spanish MICINN grants AYA2010-15685 and MEC grants AYA2010-15685, AYA2008-04211-C02-C01.

REFERENCES

[1] J. F. steffensen, "Remarks on iteration", Skand Aktuar Tidsr, vol. 16, pp. 64-72, 1933.
[2] M.A. Hernndez, Second-derivative-free variant of the Chebyshev method for nonlinear equations, J. Optim. Theory Appl., 104 (3) (2000), pp. 501?515.
[3] M.A. Hernndez, Chebyshev?s approximation algorithms, applications, Comput. Math. Appl., 4 (2001), pp. 433?445.
[4] J. Kou, Y. Li, X. Wang, A uniparametric Chebyshev-type method free from second derivatives, Appl. Math. Comput., 179 (2006), pp. 296?300.
[5] Q.B. Wu, Y.Q. Zhao, The convergence theorem for a family deformed Chebyshev method in Banach space, Appl. Math. Comput., 182 (2006), pp. 1369?1376.
[6] M. Frontini, E. Sormani, Third-order methods from quadrature formulae for solving systems of nonlinear equations, Appl. Math. Comput., 149 (2004), pp. 771?782.
[7] J. Kou, Y. Li, On Chebyshev-type methods free from second derivative, Comm. Numer. Method Eng., 24 (2008), pp. 1219?1225.
[8] J. Kou, Y. Li, Modified Chebyshev?s method free from second derivative for non-linear equations, Appl. Math. Comput., 187 (2) (2007), pp. 1027?1032.
[9] J.R. Sharma, R.K. Guha, A family of modified Ostrowski methods with accelerated sixth-order convergence, Appl. Math. Comput., 190 (2007), pp. 111?115.
[10] B. Neta, A sixth-order family of methods for nonlinear equations, Int. J. Comput. Math., 7 (1979), pp. 157?161.
[11] C. Chun, B. Neta, A new sixth-order scheme for nonlinear equations, Appl. Math. Lett. (2011), doi:10.1016/j.aml.2011.08.012.
[12] C. Chun, Y. Ham, Some sixth-order variants of Ostrowski root-finding methods, Appl. Math. Comput., 193 (2003), pp. 389?394.
[13] H. T. Kung, J. F. Traub: Optimal order of one-point and multipoint iteration. Journal of the ACM, 21 (1974), 643?651.
[14] A. M. Ostrowski: Solution of Equations and Systems of Equations, Academic Press, New York-London, 1966.
[15] J. F. Traub: Iterative Methods for the Solution of Equations, Prentice Hall, New York, 1964.
[16] M. Grau, J.L. Diaz-Barrero, An improvment of Ostrowski root-finiding method, J. Math. Anal. Appl., 173 (2006) 450-456.
[17] Alicia Cordero and Jos L. Hueso and Eulalia Martnez and Juan R. Torregrosa, A family of iterative methods with sixth and seventh order convergence for nonlinear equations, J. Mathematical and Computer Modelling, 52 (2010) 1490-1496.
[18] Sanjay K. Khattri, Joannis K. Argyros, Sixth order derivative free family of iterative methods, J. Appl. Math. Comput., 217 (2011), 5500-5507
[19] R. Thukral, New family of higher order Steffensen-type methods for solving nonlinear equations, J. of Mod. Meth. in Numer. Math., Vol. 3, No. 1, 2012, 1-10
[20] F. Soleymani, V. Hosseinabadi, New third- and tixth-order derivative-free techniques for nonlinear equations, J. of Math. Research, Vol. 3 N0. 2; May 2011.
[21] Sanjay Khattri, Ioannis Argyros, Unification of sixth-order iterative methods, J. of Mathematical Sciences 2013, 7:5, doi:10.1186/2251-7456-7-5.
[22] F. Soleymani, Efficient Sixth-Order Nonlinear Equation Solvers Free from Derivative, World Applied Sciences Journal 13 (12): 2503-2508, 2011.
[23] http://www.maplesoft.com/products/maple/
[24] Laila M. Assas et al., Eighth-order Derivative-Free Family of Iterative Methods for Nonlinear Equations, J. of Mod. Meth. in Numer. Math. (2013), in press.
[25] R. Thukral. A family of three-point derivative-free methods of eight-order for solving nonlinear equations. J. of Mod. Meth. in Numer. Math., 3(2):11-21, 2012.
[26] Fazlollah Soleymani, Some efficient seventh-order derivative-free families in root-finding, Opuscula Math. 33, no. 1 (2013), 163?173, http://dx.doi.org/10.7494/OpMath.2013.33.1.163

Dept. de Fsica i Enginyeria Nuclear, Universitat Politcnica de Catalunya, Comte d'Urgell 187, 08036 Barcelona, Spain

E-mail address: `fayyaz.ahmad@upc.edu, domingo.garcia@upc.edu`